

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/361869117>

Fate and impact of nano/microplastic in the geoenvironment — ecotoxicological perspective

Article in *Journal of Environmental Geotechnics* · September 2022

DOI: 10.1680/jenge.22.00053

CITATIONS

0

READS

328

8 authors, including:



Brendan C. O'Kelly
Trinity College Dublin

216 PUBLICATIONS 3,608 CITATIONS

[SEE PROFILE](#)



Olga Pantos
Institute of Environmental Science and Research

65 PUBLICATIONS 3,650 CITATIONS

[SEE PROFILE](#)



Theo S. Sarris
Environmental Science & Research

44 PUBLICATIONS 174 CITATIONS

[SEE PROFILE](#)



Venkata Siva Naga Sai Goli
Indian Institute of Technology Bombay

27 PUBLICATIONS 349 CITATIONS

[SEE PROFILE](#)

Some of the authors of this publication are also working on these related projects:



Phyllites [View project](#)



A Study on Ground Water Quality Analysis in AP CRDA Region [View project](#)

Cite this article

O'Kelly BC, Pantos O, Weaver L *et al.*
Fate and impact of nano/microplastic in the geoenvironment — ecotoxicological perspective.
Environmental Geotechnics,
<https://doi.org/10.1680/jenge.22.00053>

Research Article

Paper 2200053

Received 08/04/2022; Accepted 01/07/2022

ICE Publishing: All rights reserved

Fate and impact of nano/microplastic in the geoenvironment — ecotoxicological perspective

Brendan C. O'Kelly MEngSc, PhD, FTCD

Associate Professor, Department of Civil, Structural and Environmental Engineering, Trinity College Dublin, Dublin, Ireland (Orcid:0000-0002-1343-4428) (corresponding author: bokelly@tcd.ie)

Olga Pantos PhD

Senior Scientist, Health and Environment Division, Institute of Environmental Science and Research, Christchurch Science Centre, Christchurch, New Zealand (Orcid:0000-0001-5454-9056)

Louise Weaver PhD

Science Leader, Health and Environment Division, Institute of Environmental Science and Research, Christchurch Science Centre, Christchurch, New Zealand (Orcid:0000-0002-3750-868X)

Theo S. Sarris MEng, MSc, PhD, CPEng

Science Leader, Health and Environment Division, Institute of Environmental Science and Research, Christchurch Science Centre, Christchurch, New Zealand (Orcid:0000-0002-1429-2185)

Venkata Siva Naga Sai Goli MTech, SM ASCE, SM ASTM

Research Scholar, Department of Civil Engineering, Indian Institute of Technology Bombay, Mumbai, India (Orcid:0000-0003-1916-725X)

Arif Mohammad PhD

Research Associate, School of Engineering, Cardiff University, Cardiff, UK; Formerly Post-doctoral Research Fellow, Department of Civil Engineering, Indian Institute of Technology Bombay, Mumbai, India

Prithvendra Singh MTech, SM ASCE, SM ASTM, SM ICE (UK)

Research Scholar, Department of Civil Engineering, Indian Institute of Technology Bombay, Mumbai, India (Orcid:0000-0001-9979-7694)

Devendra Narain Singh PhD, FASCE, FNAE, FICE

DL Shah Chair Professor for Innovation, Department of Civil Engineering, Indian Institute of Technology Bombay, Mumbai, India (Orcid:0000-0003-3832-1507)



Plastic pollution in the terrestrial environment is emerging as another significant human-made threat to ecosystem function and health. Plastic contamination can range from the macro- to the nanoscale, and environmental impacts are evident at each level. Although significant knowledge gaps remain regarding the interactions between the natural environment and nano- and microplastics (NMPs), there is an increasing body of evidence concerning detrimental effects on a wide range of taxa. The surface properties of NMPs lead to the adsorption of heavy metals, endocrine-disrupting chemicals, antibiotics and other persistent organic pollutants, which, therefore, can result in their co-migration in the terrestrial environment. Although airborne and dietary transmission routes of NMPs have been observed, their effects on human health are still not fully understood, which is of concern to the scientific community. This state-of-the-art review paper firstly examines available evidence for, and knowledge of, various sources of NMP contamination to the terrestrial environment. Attention then focuses on (a) the biological processes from the source to soils and plants, (b) potential impacts of NMPs on soil and subsurface ecosystems, (c) trophic interactions and function and (d) implications for environmental and human health. This paper concludes by identifying knowledge gaps and presents recommendations on prioritised research needs.

Keywords: geoenvironment/land contamination/pollution migration/control/public health/UN SDG 3: Good health and well-being/UN SDG 12: Responsible consumption and production/UN SDG 15: Life on land/waste management & disposal

Introduction

All life on earth, from microorganisms to humans, is becoming increasingly exposed to plastic pollution. Society's addiction to plastics, from uses in clothing to food packaging, is driving a spiral of destruction of the planetary ecosystem on a scale comparable with that of climate change, while simultaneously being intricately linked with it. The impact of plastic pollution in oceans, on coastlines, in soils and in freshwater environments, and ultimately in food, water and air, has become a hot topic worldwide, and increasingly for public health specialists. While the focus so far has predominantly been on the levels, types and impacts of plastics in marine environments, it is estimated that the levels of plastics entering terrestrial ecosystems are between 4 and 23 times greater than those

entering the oceans (Horton *et al.*, 2017). The severity has heightened with the emergence of the coronavirus disease 2019 pandemic, which has increased single-use plastic usage globally for personal protection and hygiene and intensified the indiscriminate disposal of plastic wastes (Vaverková *et al.*, 2021).

Although no internationally agreed definition of their size range exists (Hartmann *et al.*, 2019), microplastics (MPs) are generally defined as plastic debris with particle sizes ranging between 100 nm and 5 mm (Galgani *et al.*, 2013; Thompson *et al.*, 2004), whereas plastic particles < 100 nm sized are referred to as nanoplastics (NPs) (Alimi *et al.*, 2018; Jahnke *et al.*, 2017). While there remain significant knowledge gaps around the interactions between these

Offprint provided courtesy of www.icevirtuallibrary.com
Author copy for personal use, not for distribution

plastics and the natural environment and their impacts on ecosystem function and services, it is agreed that nano- and microplastics (NMPs) have detrimental effects on a broad range of species at different trophic levels, in both aquatic and terrestrial environments, including those providing essential ecosystem services, such as earthworms (Browne *et al.*, 2013; Lwanga *et al.*, 2016). These effects, similar to those evident in marine ecosystems, may be either direct or indirect and are dependent on different characteristics, such as particle size and surface properties, of the plastic particles themselves (de Souza Machado *et al.*, 2018). For example, the surface properties of NMPs (e.g. hydrophobicity, high specific surface area and morphology) lead to adsorption of heavy metals, endocrine-disrupting chemicals, antibiotics and other persistent organic pollutants (Su *et al.*, 2021) and their co-migration in the terrestrial environment (Qi *et al.*, 2020a). Migration of NMPs through the soil can occur through numerous mechanisms, including by the ingestion and egestion of soil organisms (e.g. earthworms), translocation by water through burrows and tubes and tillage or soil wetting/drying cycles. Although understanding of the effects on human health is still in its infancy, the recent

confirmation of plastics in human blood (Leslie *et al.*, 2022) is of significant concern, and it is important that routes of transfer are identified to mitigate exposure. For example, NMPs ingested by soil organisms (e.g. earthworms) can enter the human food chain through predators, such as commercial poultry (chicken) (Wang *et al.*, 2020). Trophic transfer within the human food chain is also suggested to occur through the consumption of plants, with recent studies suggesting the transfer of NMPs from the soil into the edible parts of plants (roots, leaves and fruits) (Oliveri Conti *et al.*, 2020; Schwab *et al.*, 2020). A secondary route of exposure of NMPs from soil to humans through the consumption of plants may occur due to aeolian NMPs becoming deposited on leaves (Liu *et al.*, 2020) or through consumption of livestock that have fed on aeolian-NMP-contaminated leaves.

Using the Scopus search engine, research-article-, review-, conference-paper- and book-chapter-type papers published in the last 10 years were screened using five different keyword combinations: (a) 'MPs in oceans', (b) 'MPs in marine', (c) 'MPs in seawater', (d) 'MPs in terrestrial' and (e) 'MPs ecotoxicological




Figure 1. Number of publications on various MP-related topics published in the last 10 years (up to June 2022): (a) 'MPs in oceans'; (b) 'MPs in marine'; (c) 'MPs in seawater'; (d) 'MPs in terrestrial'; (e) 'MPs ecotoxicological effects'

Offprint provided courtesy of www.icevirtuallibrary.com
Author copy for personal use, not for distribution

effects'. When plotted (Figure 1), the synthesis of data shows a dramatic increase in the numbers of publications for each category produced on a yearly basis. Also, it can be realised that, compared with the other categories, the research works on MPs considering the marine environment are much greater, implying the current focus of researchers on MP studies. Therefore, MPs in terrestrial areas also need to be explored, to obtain a better perspective of their presence in all environments. Furthermore, from a total of 32 publications available for the keyword 'MPs ecotoxicological in terrestrial', only 13 are review-type papers, with very few of them concentrated on the ecotoxicological effects of NMPs on the terrestrial ecosystem. Hence, it necessitates a comprehensive review to analyse critically the existing literature, as done in the present paper, with a leaning on environmental geotechnics aspects.

This state-of-the-art review paper begins with an appraisal of available evidence for, and knowledge of, various sources of NMP contamination to the terrestrial environment. The focus then turns to NMP transport in the terrestrial environment; its impacts on soil properties, soil biology and subsurface ecosystems; and potential implications for human health. The paper concludes by identifying knowledge gaps in the literature and presents recommendations on prioritised research needs.

NMP sources and composition

The types and composition of NMPs entering the environment can vary significantly between the upstream sources from which they originate. Dominant sources of NMPs to soils include industry, domestic households, vehicular transport, wastewater treatment plants (WWTPs), agriculture, horticulture and landfills (Figure 2). NMPs can be divided into two categories, primary and secondary. Primary NMPs are manufactured, and include, for example, microbeads for use in personal care products (Golwala *et al.*, 2021). Secondary NMPs are formed from fragmentation of plastics through environmental, chemical and (or) biological processes, including, for instance, microfibres arising from synthetic textiles. The morphotype of NMPs (fibre, fragment, bead or film) plays an important role in determining their fate and transport in the geoenvironment, as well as their impact on organisms that they interact with (Ren *et al.*, 2021; Shamskhany *et al.*, 2021).

WWTP practices have a significant role in the distribution of NMPs into the environment. Dominant morphotypes of plastic found in municipal wastewater are synthetic fibres, followed by fragments and films, and in some locations microbeads, when they are still permitted for use in personal care and domestic




Figure 2. NMP transport and fate in relation to soil health

Offprint provided courtesy of www.icevirtuallibrary.com
Author copy for personal use, not for distribution

cleaning products (Golwala *et al.*, 2021; Rasmussen *et al.*, 2021; Ruffell *et al.*, 2021). As well as being represented by different morphotypes, the MPs in wastewater consist of a broad range of polymer types, the dominant ones being polyester (PEST), polyethylene (PE) and polypropylene (PP) (Ruffell *et al.*, 2021). WWTPs can achieve a high MP removal efficiency (80–99%), although much is physical removal, such that a large percentage of the MPs end up in the sludge residue (Carr *et al.*, 2016; Rasmussen *et al.*, 2021). In the study by Rasmussen *et al.* (2021), a mass balance was performed for a large municipal WWTP in Sweden (receiving 201.2 kg/day of plastic waste). They concluded that most plastics were removed at the screens entering the WWTP (the 20 and 2 mm bar screens respectively removed 38.2% (equivalent to 76.8 kg/day) and 35.2% (equivalent to 70.8 kg/day) of the plastics), with 13.6% (equivalent to 27.3 kg/day) contained in the residual sludge. Only a small proportion (0.7 kg/day) was contained in the WWTP effluent. However, it should be noted that there was a large proportion missing from the mass balance (12.7% or 25.5 kg/day), attributed to sampling and analysis errors and also potential loss in the digesters and activated sludge process. The high removal by the screens emphasises the importance of adequate management of screen waste. In countries where screened waste is incinerated, the potential for NMP release into the environment is small. However, many countries dispose of screened waste in landfills, which may have a much bigger impact on the environment through transport into groundwater. The fact that most NMPs end up in sludge or biosolids has large implications in the terrestrial environment, since biosolids are applied to land as soil conditioners and fertilisers. It has been estimated that the quantity of MPs annually applied to farmland in Europe through addition of sludge could be as high as 40 000–50 000 t, and between 63 000 and 430 000 t in North America (Nizzetto *et al.*, 2016). More recent studies confirmed these numbers, with up to 19 000 t of MPs annually applied to agricultural soils in Australia (Ng *et al.*, 2018).

There is a lack of research on the extent of NMP release through on-site wastewater treatment systems – for example, septic tank systems and package plants. It is logical that a concentration of NMPs occurs in these systems since they are not designed to remove plastic waste. Consequently, NMPs end up entering the environment through the disposal or land application field or through sludge removal. Disposal fields, normally comprising sand or fine gravel, are used to discharge effluent in a controlled manner into the subsurface, such that any NMPs present will also be discharged into the disposal field. Effluent passes through the disposal field before entering the vadose (unsaturated) zone and then groundwater. The design of disposal fields is such that additional physical removal is achieved before reaching the groundwater. Removal largely depends on the size and surface charge of the NMP particles.

More research is needed to ascertain the NMP removal rates through WWTP and on-site wastewater treatment systems.

If NMPs are removed and retained in the sludge, they normally (re-)enter the municipal wastewater treatment system, where the potential for their removal has been explained earlier.

Another source of NMPs is food production farms, where plastics are widely used, from netting to plastic mulch film. Although designed to withstand environmental conditions through incorporation of additives (e.g. ultraviolet (UV) stabilisers) in their manufacturing process, plastics degrade over time because of weathering and mechanical wear and tear, forming fragments (i.e. NMPs) that stay in the soil (Kasirajan and Ngouajio, 2012; Steinmetz *et al.*, 2016). In some locations, it is common practice that, rather than being collected and removed after the growing season, horticultural plastics (e.g. mulch film) are ploughed into the soil. Other sources of plastics in soils are the application of organic and inorganic fertilisers. For instance, many pelletised synthetic fertilisers possess polymer-based microcapsules (Katsumi *et al.*, 2021). After their gradual breakdown, allowing slow release of the fertiliser, the synthetic polymer remains in the soils as NMPs. The application of natural organic fertilisers resulting from the recycling of municipal green waste (either compost or anaerobic digestate) also represents a significant source of plastic particles to soils because of the high levels of plastics that find their way into the food waste (Golwala *et al.*, 2021; Weithmann *et al.*, 2018). Although the role of agricultural and horticultural practices in the contribution of NMPs in soils has been identified, their levels are not fully understood and further research is needed to establish associated risks.

There is evidence that landfills are defused sources of NMPs to their immediate environment (Table 1), being identified in the surrounding air, groundwater, surface water and soil (O'Kelly *et al.*, 2021). Unmanaged, old or legacy landfills do not include protective liners, covers and leachate and gas collection systems, such that leachate can migrate into the environment. The leachate contains high levels of primary and secondary MPs due to mechanical fragmentation of larger plastic items, the presence of discarded products (including personal care and industrial cleaning products) and NMPs present in landfilled WWTP sludge (Guerranti *et al.*, 2019; Sobhani *et al.*, 2020). Dominant polymers found in leachate are PE and PP (see Table 1), accounting for almost 99.4% of NMPs present, mainly because (a) the majority of the municipal plastic waste is derived from either PE or PP (Goli *et al.*, 2020; He *et al.*, 2019) due to their high usage levels and (b) since the density of these polymers is $<1.0 \text{ g/cm}^3$, they float and are transported by the leachates. The levels of MPs in solid refuse samples have been found to be significantly higher ($62\,000 \pm 23\,000 \text{ particles/kg}$ (Su *et al.*, 2019)) than those found in sewage sludge (between 4196 and 15 385 particles/kg (Mahon *et al.*, 2017)). It should be kept in view that the physico-chemical attributes of NMPs generated in municipal solid waste facilities would be distinctive, due to the presence of a mixture of both conventional and biodegradable polymers.

Vehicular transport also acts as a significant source of plastics to the environment (Schwarz *et al.*, 2019), in the form of brake-wear

Offprint provided courtesy of www.icevirtuallibrary.com
Author copy for personal use, not for distribution

Table 1. Concentrations of plastic particles identified in different components of landfills and their surrounding ecosystems

Source/ ecosystem component	Location	Size: μm	Average MP concentration	Salient observations	Reference
Leachate	Shanghai, Wuxi, Suzhou and Changzhou, China	100–5000	0.42–24.58 ^a	PE and PP were dominant MPs 99.36% of MPs were derived from fragmentation of plastic waste buried in landfills 77.48% of MPs sized between 0.1 and 1.0 mm	He <i>et al.</i> (2019)
	Shanghai, China	20–5000	8 \pm 3 (4–13) ^a	Dominant morphotype of MPs was fibres (60%) and film Average MP concentrations in young (<3 years), medium (~10 years) and old leachates (>20 years) were 8, 10 and 4 MP particles/l, respectively Observed nine different polymer types present based on functional groups identified using FTIR spectroscopy	Su <i>et al.</i> (2019)
Refuse	Shanghai, China	20–5000	62 000 \pm 23 000 (20 000–91 000) ^b	MP abundance in young (<3 years), medium (~10 years) and old refuse (>20 years) of 83 000 \pm 10 000, 68 000 \pm 6000 and 36 000 \pm 14 000 particles/kg, respectively Observed 15 different types of thermoplastic and thermoset polymers present Most dominant shape of MPs was fibres (59.82%)	Su <i>et al.</i> (2019)
Compost	Paris, France	0.45–5000 ^c	^d	Used pyrolysis–gas chromatography–mass spectroscopy for detection of MP presence in soils	Watteau <i>et al.</i> (2018)
Groundwater	Chennai, India	0.45–5000	Perungudi site: 33 (7–80) ^a Kodungaiyur site: 12 (3–23) ^a ^e	90% of MPs were derived from buried plastics Major polymer types in groundwater were PS and PP Dominant colours of MPs were white (38%), black (27%), red (18%), green (8%) and blue (6%)	Manikanda Bharath <i>et al.</i> (2021)
Soil	Dhaka, Bangladesh	1–2000		Samples were collected at two different depths: topsoil and 0–20 cm layer	Afrin <i>et al.</i> (2020)
Lichen	Tuscany, Italy	<5000	Close, 79 000 (0–95 000) ^b Intermediate (i.e. 200 m distant), 13 000 (0–15 000) ^b Remote (i.e. 1500 m distant), 7000 (3000–9000) ^b	Influence of landfill on MP concentration in lichen was determined; MP concentration reduced with increasing distance from landfill.	Loppi <i>et al.</i> (2021)

^a MP particles/l

^b MP particles/kg dry mass of sample/matrix tested

^c Lower and upper boundaries of size range decided based on pore size of filter paper and aperture size of sieve, respectively, used in the experiments

^d Quantification cannot be obtained since, in this method, the MPs are degraded at higher temperatures before the detection stage, using mass spectrometry

^e Not mentioned in the original paper, as the detection technique of FTIR spectroscopy, using the potassium bromide (KBr) pellet method, cannot be used to quantify the MPs

FTIR, Fourier transform infrared; PE, polyethylene; PP, polypropylene; PS, polystyrene

and tyre-wear particles (Evangelidou *et al.*, 2020). The wearing process depends on the vehicle and pavement characteristics and type of tyre (Grigoratos and Martini, 2014). Kole *et al.* (2017) reported movement of break- and tyre-wear particles owing to washout and run-off to freshwater and marine ecosystems. The size of these particles can be <10 μm , and thus, they can also remain airborne for longer periods (Harrison *et al.*, 2012). Apart from these sources, polymerised bitumen is another supply of NMPs (Rødland *et al.*, 2022).

Other potential sources of local MP contamination of soils and groundwater are the addition of synthetic polymer-based (usually PP) fibres, as strength enhancement (reinforcement) for ground

improvement, and synthetic polymers, including tyre derived aggregates (TDAs) used, for instance, as partial replacements for natural soils and aggregates in the construction of road embankments and pavement subgrades and as lightweight backfill to bridge abutments and retaining walls. As well as the MP particles derived from weathering and physical breakdown of the TDA additive in situ, there is the potential toxicity of leachates containing heavy metals and other chemicals common in TDA materials, as additional sources of contamination to the terrestrial environment (Váverková and O'Kelly, 2022). For instance, Tallec *et al.* (2022) emphasised that the use of rubber-based products (e.g. crumb rubber granulates) can induce 'rubber contamination' by releasing micro-rubber and (or) constitutive compounds (added during the tyre manufacturing processes) that

Offprint provided courtesy of www.icevirtuallibrary.com
Author copy for personal use, not for distribution

leach out by the action of water. From the studies of Šourková *et al.* (2021a, 2021b), it is evident that leachates from waste tyre fractions are phytotoxic to highly phytotoxic to higher plants.

Further geotechnical engineering examples include polystyrene (PS)-based lightweight engineering fills (Abbasimaedeh *et al.*, 2021; O’Kelly and Soltani, 2022a) or use of dredge sediments from water resources as fill materials (Monkul and Özhan, 2021). Dredged sediments may contain significant amounts of MPs owing to their free dispersion in the aquatic environment (Ji *et al.*, 2021). Synthetic polymer-based fibres are also used as secondary additives to increase the ductility of stabilised soils that are mainly used as barriers in landfills or subgrade soils. Expanded polystyrene (EPS), as blocks (i.e. geofoam) or as myriads of discrete beads mixed with soil in situ, is a preferred lightweight fill for embankment construction over soft soil deposits, as retaining wall backfill and to protect culverts and buried pipelines. At the end of life of earth structures constructed using soil–EPS bead mixtures or using myriads of discrete plastic fibres mixed randomly with soil in situ, the EPS beads and plastic fibres contained therein are not alienable from the soil. Hence, they are not recyclable (Abbasimaedeh *et al.*, 2021; O’Kelly and Soltani, 2022a, 2022b) and have the potential to cause substantial local NMP contamination of soil and groundwater. Based on environmental concerns and also due to the substantial deterioration in geomechanical behaviour/properties for increasing EPS additive content, Abbasimaedeh *et al.* (2021) went so far as to recommend that geotechnical engineering practice discontinue the approach of adding particulate EPS beads to soil for producing uncemented lightened fills.

Note that even though NPs and small MPs are widely present in the environment, they are often not detected or accurately quantified in environmental matrices, including soils, because of current methodological and analytical limitations – identified as a major shortcoming for present research efforts (Goli *et al.*, 2022; O’Kelly *et al.*, 2021).

NMP transport in terrestrial environment

While NMP transport above ground is driven by wind and surface water movement, once NMPs enter the soil matrix, their horizontal and vertical migration are controlled by soil physical properties, soil biota, agronomic practices and hydrological conditions (i.e. rainfall frequency and intensity) (Rillig *et al.*, 2017a). Due to the pore sizes of fine-grained soils being smaller than the size of many NMPs, a large proportion of NMPs are retained by the upper soil layer (Ren *et al.*, 2021). However, owing to the absence of UV light and low stable temperatures, once in the subsurface, NMPs can accumulate and remain unaltered for prolonged periods (Otake *et al.*, 1995). Some studies have indicated that not only are soils NMP sinks, but they can also provide a feasible transport entryway to subsurface receptors through advective or colloidal transport. For instance, O’Connor *et al.* (2019) found that MPs can undergo significant vertical migration in sandy soils, with transport distances increasing for reducing MP particle size and (or) under wetting/drying cycles. Similar transport characteristics should be expected in soils with high proportions of macropores (e.g. bio-pore-rich loamy soils, organic-rich soils, leptosols and vertisols), for which preferential flow and transport can occur (Bläsing and




Figure 3. Schematic representation of NMP effects and impacts on soil health

Offprint provided courtesy of www.icevirtuallibrary.com
Author copy for personal use, not for distribution

Amelung, 2018). Vertical transport to deep soils and the vadose zone is further accentuated by bioturbation, in the form of fragmentation and size-selective transport (Huerta Lwanga *et al.*, 2017; Rillig *et al.*, 2017b; Zhu *et al.*, 2018). In this context, MPs are prone to degradation in terrestrial environments, which decreases their particle size, increases their specific surface area and oxygen-containing functional groups and enhances the potential chance of attachment (i.e. through sorption, electrostatic force, etc.) of microorganisms, heavy metals and other pollutants present in soil (Golwala *et al.*, 2021).

Impacts on the geoenvironment

Once NMPs arrive on soils, one must consider the role of environmental conditions affecting their fate and potential impacts (see Figure 3). For instance, weathering due to UV radiation and hydrolysis affects the physical characteristics of plastics, including polymer structure changes, alteration of the surface texture and promotion of fragmentation. Further, when considering NMP impacts on the terrestrial environment, one cannot only consider the physical NMPs themselves but must also consider their associated chemicals – that is, in the manufacturing of plastics, various chemical additives, ranging from plasticisers to UV stabilisers, are included to suit the end-product use. This section of the paper focuses on NMP impacts on the soil properties, soil biology and subsurface ecosystems.

Soil properties

The impact of NMPs on the physical structure of the soil is beginning to be realised, particularly in studies from China, where, since the late 1970s, there has been an increasingly common practice of plastic film mulching (PFM) for improving cash-crop yields (Huang *et al.*, 2020; Ng *et al.*, 2018; Qi *et al.*, 2020b). Studies have shown that PFM can have physical effects on soil ecosystems, including reduced soil porosity (Koskei *et al.*, 2021), changed air circulation and altered microbial communities (Li *et al.*, 2014; Muroi *et al.*, 2016), increased soil water-retention capacity (de Souza Machado *et al.*, 2019) and greater greenhouse gas emissions (Cuello *et al.*, 2015). Additionally, Wang *et al.* (2016) found that the practice of PFM significantly reduces soil biomass carbon and nitrogen contents, soil metabolic activity and microbial function and activity. Weathering and physical breakdown of plastic mulch film results in the formation and accumulation of NMPs within the soil matrix. NMP presence has also been shown to have a physical impact on soil mesofauna owing to the reduction in porosity, effectively causing their immobilisation (Kim and An, 2019). Luo *et al.* (2020) reported that physico-chemical factors, in particular pH and soil organic carbon, play a role in the attachment of NMPs (0.047 µm sized PS). Five soils were tested, and a strong positive correlation was seen between the NMP's sorptive capacity to the soil and both organic carbon and Fe₂O₃ soil content. In contrast, the higher the clay content, the lower the sorption of NMPs to the soils. Luo *et al.* (2020) also reported that the sorption capacity of all the test soils decreased as pH increased. They concluded that the main attractive forces contributing to NMP sorption to soils were

electrostatic interaction and hydrophobic interaction. A more recent study by Wang *et al.* (2022) agreed with these findings and added that attachment of MPs was significantly correlated with the zeta potential of soils.

Soil biology

Plastics that enter the environment will inevitably be colonised by microbial communities. In any food web, microbes are keystone organisms, often existing in biofilms to enhance survival and for protection from predation. In forming biofilms, microbes will interact with plastic in any environment, as a surface to attach to or as a supplementary carbon source. Nevertheless, synthetic plastics are recalcitrant, and few studies have demonstrated actual plastic degradation directly arising from microbial action, most of them demonstrating that the additives used in plastic manufacturing may be degraded, rather than the plastic polymer itself. Biodegradable plastics, manufactured with aliphatic PESTs (e.g. polyhydroxyalkanoate and polylactic acid), have been shown to degrade, although they may require specific conditions not present in the soils. Even though there are numerous studies demonstrating microbiomes (biofilms) associated with plastics across aquatic and terrestrial environments, the impact of plastics on these communities is sparsely investigated (Lear *et al.*, 2021; Ng *et al.*, 2021). A recent study demonstrated that distinct communities colonise MPs in soil (Zhang *et al.*, 2019a), but it did not investigate the impact of this on the soil ecosystem. Ng *et al.* (2021) investigated the impact of polyethylene terephthalate and low-density PE (LDPE) MPs by comparing with a control soil from a forest environment. Shifts were observed in microbial composition, both between plastics and relative to the control soil, indicating that the chemical composition (as different plastics age in the soil) will have an impact on the soil microbiome. Moreover, the presence of LDPE MPs resulted in a seven- to eightfold increase in carbon dioxide (CO₂) production compared with the control soil.

A direct impact of colonisation of MPs with biofilm communities is degradation or fragmentation/disintegration. There is demonstrated evidence (predominantly in a laboratory setting) that bacterial and fungal species can degrade MPs (Gambarini *et al.*, 2021; Lear *et al.*, 2021; Ng *et al.*, 2018; Wei and Zimmermann, 2017). Another consequence of colonisation is that soil detritivores consume microbial biofilms and, hence, may also incidentally engulf MPs (Guo *et al.*, 2020). Some studies have demonstrated that detritivores, such as earthworms, consume colonised organic matter (including MPs) over fresh material (Huerta Lwanga *et al.*, 2017; Rillig *et al.*, 2017a, 2017b), producing detrimental effects (Lwanga *et al.*, 2016). These include intestinal problems, blockages and stopping vital nutrients from being taken in by the animal or, alternatively, can give the animal a feeling of being full, so that they do not feed and starve as a result. However, detailed studies on the impacts of NMP ingestion by soil biota are limited. A reduction in activity in detritivores will change the soil physical properties, such as nutrient addition and bioturbation. This could reduce soil structure and bioavailability of nutrients for plant growth. More studies are needed

Offprint provided courtesy of www.icevirtuallibrary.com
Author copy for personal use, not for distribution

in this area to pinpoint exact impacts across biological function and ecosystem services.

Subsoil transport

Although soils may be predominantly a sink for plastics, it is inevitable that degradation in solids will lead to NMPs becoming a source of NMPs in groundwater environments. Physical processes, such as described in the section headed 'NMP transport in terrestrial environment', can lead to contamination of the vadose zone and ultimately the groundwater environment. Recent studies have begun to illustrate the risk to groundwater from landfill sites and agricultural soils (Lwanga *et al.*, 2022; Manikanda Bharath *et al.*, 2021; Ren *et al.*, 2021; Samandra *et al.*, 2022; Zhang *et al.*, 2022). Once in the groundwater environment, NMPs can be transported in significant quantities in alluvial aquifers (Goeppert and Goldscheider, 2021). There is a sparsity of other studies looking at the transport and fate of NMPs in groundwater (Viaroli *et al.*, 2022). The potential of NMP transport depends on the lithology and geochemical conditions, such as clay and colloidal materials present. Also, environmental factors and soil parameters (e.g. pH, primary cations and iron (Fe) mineral and organic matter) influence the transport behaviour of MPs in the soil matrix (Ren *et al.*, 2021). One of the key issues with the study of NMPs in groundwater is the lack of consistent methods for sampling and quantification (Viaroli *et al.*, 2022). To date, only five articles (Goeppert and Goldscheider, 2021; Johnson *et al.*, 2020; Panno *et al.*, 2019; Samandra *et al.*, 2022; Selvam *et al.*, 2021) present sufficient information on the groundwater sampled and the hydrogeological information needed for assessment (Viaroli *et al.*, 2022). The NMP concentrations varied by orders of magnitude in different aquifer materials, but, as expected, they followed a pattern of increasing concentration with large pore size or fractured systems present. Karstic systems are likely to have NMP-size distributions similar to those found in surface waters (Panno *et al.*, 2019). Alluvial aquifers have been shown to transport a wide range of NMP sizes, down to a 1 µm size (Goeppert and Goldscheider, 2021).

Furthermore, with very little known about groundwater ecosystems themselves, the impact on the groundwater ecosystems present is unaccounted for, such that understanding the NMP effect on their function and its subsequent impact on the water cycle is currently unknown. The fact that globally around 2 billion people depend on groundwater resources makes this a critical issue to be addressed urgently.

Ecotoxicity

Studies on the impacts and ecotoxicological effects of NMPs on eukaryotic organisms are more common than those on prokaryotes, with studies across springtails, earthworms, nematodes, arthropods, isopods and mites. In a recent investigation on the plastic additive bisphenol A, Gerhardt (2019) indicated acute and chronic toxicity effects on surface and groundwater crustaceans. Compared with detritivores, higher sensitivities were evident for filter-feeding crustaceans, as may be expected for a dissolved toxin. Gerhardt

(2019) concluded that the groundwater crustacean isopod *Proasellus slavus* was the most sensitive to both acute and chronic exposure, such that it could be a useful indicator species. Oxidative stress, histopathological changes and reproduction impediment have been indicated in response to MP exposure in the earthworm *Eisenia andrei* (Jiang *et al.*, 2020; Kwak and An, 2021; Lackmann *et al.*, 2022). Studies on predator–prey interactions at the higher levels of the food web are sparser, with most only showing indirect evidence (Helmlinger *et al.*, 2020).

Until recently, flora has been largely overlooked regarding ecotoxicological effects from plastics, those exceptions concerning the terrestrial environment including the investigations by Allouzi *et al.* (2021), Mateos-Cárdenas *et al.* (2021) and Ng *et al.* (2021). For instance, Mateos-Cárdenas *et al.* (2021) showed that plants can adsorb or internalise NMPs, with the mechanisms suggested to be mostly due to electrostatic forces or entrapment in uneven surfaces on the plant. Studies have shown that NPs can also cross membrane boundaries and enter plant cells, suggesting that a toxicological effect is possible (Azeem *et al.*, 2021; Luo *et al.*, 2022; Mateos-Cárdenas *et al.*, 2021). It has been noted that, to date, most of the studies on plant interactions and impacts have been performed in controlled laboratory environments and on single species. There is still work that is needed to assess the impact of NMPs in the environment and at an ecosystem level. In the limited studies conducted, to date, NMP effects have been reported on plant germination, elongation growth and biomass. Studies that have investigated the toxic effects of plastic particles on plant photosynthesis have varied in their findings, with some noticing no effect on photosynthetic activity in controlled experiments on single species (Dovidat *et al.*, 2020; Mateos-Cárdenas *et al.*, 2019), while others (Gao *et al.*, 2019; Qi *et al.*, 2018, 2020b) have found negative effects. This variability points to the sparsity of available data and that further studies are required on a variety of plants and plastics, investigating pertinent environmental conditions.

NMP impacts on human health

The full extent of potential impacts, particularly on humans, of exposure to NMPs remains unknown. The current agreed routes of exposure of plastic particles for humans (see Figure 4) include (a) inhalation (Amato-Lourenço *et al.*, 2020; Enyoh *et al.*, 2019), synthetic fibres having been found in lung tissue (Pauly *et al.*, 1998), and (b) ingestion of water and food (both fresh and processed) (Daniel *et al.*, 2020; Dessì *et al.*, 2021; Kedzierski *et al.*, 2020; Senathirajah *et al.*, 2021). There is currently no research supporting the ability of NMPs to penetrate the surface of the skin, but it is thought that they may enter the body through sweat glands, skin wounds or hair follicles (Schneider *et al.*, 2009).

Atmospheric NMPs have been demonstrated in multiple locations globally, with them found in the air above urban areas, as well as in remote locations. This supports the hypothesis that their global transportation occurs (Zhang *et al.*, 2019b), although the process by which aeolian NMPs evolve from soils is not yet clear. It is assumed that NMPs behave similarly to nanoparticles,

Offprint provided courtesy of www.icevirtuallibrary.com
Author copy for personal use, not for distribution




Figure 4. NMP routes of exposure to humans: the soil–air–soil cycle

microorganisms and organic material – these can be resuspended from terrestrial surfaces and transported over long distances (Griffin, 2007; Yang *et al.*, 2016). Airborne synthetic fibres have been associated with respiratory diseases in humans (Prata, 2018; Turcotte *et al.*, 2013), owing to (a) difficulties associated in clearing them from the respiratory system, (b) the potential of the plastic to interact with other organic materials and (c) the release of chemical contaminants associated with the fibres (Enyoh *et al.*, 2019). High levels of plastic fibres present within wastewater effluent (Cao *et al.*, 2020; Dyachenko *et al.*, 2017) and biosolids (Koutnik *et al.*, 2021) applied to land may, therefore, pose a human health risk through inhalation after resuspension into the air, as well as other morphotypes also present (Enyoh *et al.*, 2019; Knobloch *et al.*, 2021).

It has been estimated that children and adults may ingest as many as 100 000 MP particles each day (Mohamed Nor *et al.*, 2021), with other studies estimating an average ingestion rate of 0.1–5 g of MPs per week through various exposure pathways (Senathirajah *et al.*, 2021). Internalisation of MPs through the dietary pathway has been confirmed by evidence of their presence

within human stool and blood samples (Leslie *et al.*, 2022; Schwabl *et al.*, 2019). Animal studies have demonstrated the translocation of NMPs from the gut (Browne *et al.*, 2008; Mattsson *et al.*, 2017), although, to date, similar data have not been possible to determine this for humans.

NMPs may be introduced into food during meal preparation (Knobloch *et al.*, 2021; Zhang *et al.*, 2020), or they may already be present within fresh food items. Besides the potential health risk posed by NMPs within plant tissues when consumed, the presence of plastics within the soil facilitates the uptake of heavy metals into plants, which may pose a direct human health risk (Wang *et al.*, 2021). However, there are huge uncertainties associated with detecting, identifying and characterising different NMPs (Goli *et al.*, 2022) in food.

Although there is still a need for far more research to determine the full impacts of NMPs on human health, there is sufficient evidence available to necessitate a precautionary approach to dealing with NMPs exposure. For instance, Ragusa *et al.* (2021) found plastic particles in all three layers of the human placenta:

Offprint provided courtesy of www.icevirtuallibrary.com
 Author copy for personal use, not for distribution

the maternal side, the foetal side and the chorioamniotic membrane. A better understanding of the potential damage that NMPs cause to humans will begin to emerge only when studies unravel their complex interactions with human organs.

Direct harmful effects of NMPs may be physical (mechanical) and/or chemical (toxicological), while indirect risks are posed by the presence of microbes that may colonise the plastic surface. The potential human health risks are dependent on multiple factors, including the plastic particles' size, morphology, age, associated chemicals (inherent and acquired) and microbes and exposure pathway. These factors may work in isolation or in synergy, with some of them determined by the source of the NMPs and the land use of the soil that they have been applied to. For instance, wastewater effluent that may be applied to land contains (a) high levels of very diverse NMPs; (b) a complex cocktail of organic and inorganic chemicals used in domestic and industrial processes, pharmaceuticals and personal care products (e.g. agrichemicals, antibiotics, biocides, UV sunscreens (Deblonde *et al.*, 2011)); and (c) a wide range of microorganisms, including potential human pathogens (Hansen *et al.*, 2018).

NMPs may pose direct physiological and microbial risks, together with a direct or indirect chemical risk. Within the soil matrix, associated chemicals, whether additives or acquired from the environment, may leach from the plastics, become mobile and be taken up by plants and therefore pose a potential health risk through ingestion (Al-Lihaibi *et al.*, 2019), or their presence may facilitate the uptake of non-NMP associated contaminants (Wang *et al.*, 2021). Many of these chemicals have already been determined to pose toxicological effects in humans and animals. For instance, common additives in plastics (e.g. phthalates, bisphenol A and bisphenol S) are considered endocrine-disrupting chemicals, having been linked with reproductive and developmental disorders, including breast cancer, blood infection, early onset of puberty and genital defects (Mishra *et al.*, 2019; Ribeiro *et al.*, 2019). With an increasing body of evidence showing that NMPs themselves are being taken up by plants (Bosker *et al.*, 2019; Li *et al.*, 2021; Liu *et al.*, 2020), soils may present an exposure pathway to humans through diet. Besides to acting as a vector for associated chemicals, once inside the gastrointestinal tract, NMPs may translocate to other organs (Browne *et al.*, 2013; Powell *et al.*, 2010), causing physiological effects. For example, the accumulation of NMPs in the liver and kidneys causes disturbance of energy and lipid metabolism and oxidative stress (Deng and Zhang, 2019).

Although not fully understood, initial reports are identifying the enrichment of antimicrobial resistance genes (ARGs) and potential human pathogens on NMPs (Shi *et al.*, 2021) present in the leachate of landfills that accept high volumes of municipal biosolid waste. The pathogens that were found in the leachate and MP environment were *Ochrobactrum anthropi*, *Acinetobacter lwoffii*, *A. baumannii*, *Afipia broomeae*, *Pseudomonas aeruginosa*, *Escherichia coli*, *Bacillus anthracis*, *Serratia marcescens* and *Aeromonas hydrophila*. These are opportunistic human pathogens, which are linked to

ARGs, and are responsible for nosocomial infection, such as bacteraemia, secondary meningitis, urinary tract infection and pneumonia. The high NMP load in wastewater effluent and biosolids, therefore, presents a potential high risk to microbiological human health, should they become associated with either food or water, or through direct contact of the contaminating soils, resulting in accidental ingestion. These changes to microbial communities may also cause significant changes in soil and subsoil ecosystem health.

Concluding remarks and way forward

In this paper, the authors reviewed available evidence for, and knowledge of, sources of plastic contamination to the terrestrial environment, its effects on key soil ecosystem functions and the potential for soil NMPs to enter the food chain through fauna and flora. Through the review, the following research gaps in the literature, whose investigation will require concerted multidisciplinary effort, are identified.

- Understanding NMP interactions with plants is still lacking, particularly at the ecosystem level, with very few studies reporting on multiple species and environmental impacts – those that have been done show varied responses.
- More studies are needed on the interaction of microbial biofilms in the soil environment and the potentials for pathogen survival and transport.
- Studies on the whole soil ecosystem remain somewhat lacking, particularly trophic cascade implications in terrestrial environments.

Although NMPs are widely present in the environment, they are often not detected or accurately quantified in environmental matrices (including soils) due to current methodological and analytical limitations. This has been identified as a major shortcoming for present research efforts in quantifying the extent and amount of NMP contamination in soils and groundwater.

There is also an urgent need to develop efficient remediation methods to improve the overall soil health. To date, efforts have concentrated on NMP reduction at the source, which appears to be the most viable and efficient way to manage the risk, control the effects and limit further spreading of NMPs. Remediation of water and soil from NMP pollution is still at very early stages, with microbial biodegradation and bioremediation of certain plastic pollutants showing promise at the experimental stage.

REFERENCES

- Abbasimaedeh P, Ghanbari A, O'Kelly BC, Tavanafar M and Irdmoosa KG (2021) Geomechanical behaviour of uncemented expanded polystyrene (EPS) beads–clayey soil mixtures as lightweight fill. *Geotechnics* **1**(1): 38–58, <https://doi.org/10.3390/geotechnics1010003>.
- Afrin S, Uddin MK and Rahman MM (2020) Microplastics contamination in the soil from urban landfill site, Dhaka, Bangladesh. *Heliyon* **6**(11): article e05572, <https://doi.org/10.1016/j.heliyon.2020.e05572>.
- Al-Lihaibi S, Al-Mehmadi A, Alarif WM *et al.* (2019) Microplastics in sediments and fish from the Red Sea coast at Jeddah (Saudi Arabia). *Environmental Chemistry* **16**(8): 641–650, <https://doi.org/10.1071/EN19113>.

Offprint provided courtesy of www.icevirtuallibrary.com
 Author copy for personal use, not for distribution

- Alimi OS, Budarz JF, Hernandez LM and Tufenkji N (2018) Microplastics and nanoplastics in aquatic environments: aggregation, deposition, and enhanced contaminant transport. *Environmental Science and Technology* **52**(4): 1704–1724, <https://doi.org/10.1021/acs.est.7b05559>.
- Allouzi MMA, Tang DYY, Chew KW *et al.* (2021) Micro (nano) plastic pollution: the ecological influence on soil–plant system and human health. *Science of the Total Environment* **788**: article 147815, <https://doi.org/10.1016/j.scitotenv.2021.147815>.
- Amato-Lourenço LF, Galvão LDS, de Weger LA *et al.* (2020) An emerging class of air pollutants: potential effects of microplastics to respiratory human health? *Science of the Total Environment* **749**: article 141676, <https://doi.org/10.1016/j.scitotenv.2020.141676>.
- Azeem I, Adeel M, Ahmad MA *et al.* (2021) Uptake and accumulation of nano/microplastics in plants: a critical review. *Nanomaterials* **11**(11): article 2935, <https://doi.org/10.3390/nano11112935>.
- Bläsing M and Amelung W (2018) Plastics in soil: analytical methods and possible sources. *Science of the Total Environment* **612**: 422–435, <https://doi.org/10.1016/j.scitotenv.2017.08.086>.
- Bosker T, Bouwman LJ, Brun NR, Behrens P and Vijver MG (2019) Microplastics accumulate on pores in seed capsule and delay germination and root growth of the terrestrial vascular plant *Lepidium sativum*. *Chemosphere* **226**: 774–781, <https://doi.org/10.1016/j.chemosphere.2019.03.163>.
- Browne MA, Dissanayake A, Galloway TS, Lowe DM and Thompson RC (2008) Ingested microscopic plastic translocates to the circulatory system of the mussel, *Mytilus edulis* (L.). *Environmental Science and Technology* **42**(13): 5026–5031, <https://doi.org/10.1021/es800249a>.
- Browne MA, Niven SJ, Galloway TS, Rowland SJ and Thompson RC (2013) Microplastic moves pollutants and additives to worms, reducing functions linked to health and biodiversity. *Current Biology* **23**(23): 2388–2392, <https://doi.org/10.1016/j.cub.2013.10.012>.
- Cao Y, Wang Q, Ruan Y *et al.* (2020) Intra-day microplastic variations in wastewater: a case study of a sewage treatment plant in Hong Kong. *Marine Pollution Bulletin* **160**: article 111535, <https://doi.org/10.1016/j.marpolbul.2020.111535>.
- Carr SA, Liu J and Tesoro AG (2016) Transport and fate of microplastic particles in wastewater treatment plants. *Water Research* **91**: 174–182, <https://doi.org/10.1016/j.watres.2016.01.002>.
- Cuello JP, Hwang HY, Gutierrez J, Kim SY and Kim PJ (2015) Impact of plastic film mulching on increasing greenhouse gas emissions in temperate upland soil during maize cultivation. *Applied Soil Ecology* **91**: 48–57, <https://doi.org/10.1016/j.apsoil.2015.02.007>.
- Daniel DB, Ashraf PM and Thomas SN (2020) Microplastics in the edible and inedible tissues of pelagic fishes sold for human consumption in Kerala, India. *Environmental Pollution* **266**(Part 2): article 115365, <https://doi.org/10.1016/j.envpol.2020.115365>.
- de Souza Machado AA, Kloas W, Zarfl C, Hempel S and Rillig MC (2018) Microplastics as an emerging threat to terrestrial ecosystems. *Global Change Biology* **24**(4): 1405–1416, <https://doi.org/10.1111/gcb.14020>.
- de Souza Machado AA, Lau CW, Kloas W *et al.* (2019) Microplastics can change soil properties and affect plant performance. *Environmental Science and Technology* **53**(10): 6044–6052, <https://doi.org/10.1021/acs.est.9b01339>.
- Deblonde T, Cossu-Leguille C and Hartemann P (2011) Emerging pollutants in wastewater: a review of the literature. *International Journal of Hygiene and Environmental Health* **214**(6): 442–448, <https://doi.org/10.1016/j.ijheh.2011.08.002>.
- Deng Y and Zhang Y (2019) Response to Uptake of microplastics and related health effects: a critical discussion of Deng *et al.*, Scientific reports 7: 46687, 2017. *Archives of Toxicology* **93**: 213–215, <https://doi.org/10.1007/s00204-018-2384-8>.
- Dessi C, Okoffo ED, O'Brien JW *et al.* (2021) Plastics contamination of store-bought rice. *Journal of Hazardous Materials* **416**: article 125778, <https://doi.org/10.1016/j.jhazmat.2021.125778>.
- Dovidat LC, Brinkmann BW, Vijver MG and Bosker T (2020) Plastic particles adsorb to the roots of freshwater vascular plant *Spirodela polyrrhiza* but do not impair growth. *Limnology and Oceanography Letters* **5**(1): 37–45, <https://doi.org/10.1002/lol2.10118>.
- Dyachenko A, Mitchell J and Arsem N (2017) Extraction and identification of microplastic particles from secondary wastewater treatment plant (WWTP) effluent. *Analytical Methods* **9**: 1412–1418, <https://doi.org/10.1039/c6ay02397e>.
- Enyoh CE, Verla AW, Verla EN, Ibe FC and Amaobi CE (2019) Airborne microplastics: a review study on method for analysis, occurrence, movement and risks. *Environmental Monitoring and Assessment* **191**: article 668, <https://doi.org/10.1007/s10661-019-7842-0>.
- Evangelidou N, Grythe H, Klimont Z *et al.* (2020) Atmospheric transport is a major pathway of microplastics to remote regions. *Nature Communications* **11**: article 3381, <https://doi.org/10.1038/s41467-020-17201-9>.
- Galgani F, Hanke G, Werner S and De Vrees L (2013) Marine litter within the European Marine Strategy Framework Directive. *ICES Journal of Marine Science* **70**(6): 1055–1064, <https://doi.org/10.1093/icesjms/fst122>.
- Gambarini V, Pantos O, Kingsbury JM *et al.* (2021) Phylogenetic distribution of plastic-degrading microorganisms. *mSystems* **6**(1): article e01112-20, <https://doi.org/10.1128/mSystems.01112-20>.
- Gao H, Yan C, Liu Q *et al.* (2019) Effects of plastic mulching and plastic residue on agricultural production: a meta-analysis. *Science of the Total Environment* **651**(Part 1): 484–492, <https://doi.org/10.1016/j.scitotenv.2018.09.105>.
- Gerhardt A (2019) Plastic additive bisphenol A: toxicity in surface- and groundwater crustaceans. *Journal of Toxicology and Risk Assessment* **5**(1): article 017, <https://doi.org/10.23937/2572-4061.1510017>.
- Goeppert N and Goldscheider N (2021) Experimental field evidence for transport of microplastic tracers over large distances in an alluvial aquifer. *Journal of Hazardous Materials* **408**: article 124844, <https://doi.org/10.1016/j.jhazmat.2020.124844>.
- Goli VSNs, Mohammad A and Singh DN (2020) Application of municipal plastic waste as a manmade neo-construction material: issues & wayforward. *Resources, Conservation and Recycling* **161**: article 105008, <https://doi.org/10.1016/j.resconrec.2020.105008>.
- Goli VSNs, Paleologos EK, Farid A *et al.* (2022) Extraction, characterisation and remediation of microplastics from organic solid matrices. *Environmental Geotechnics*, <https://doi.org/10.1680/jenge.21.00072>.
- Golwala H, Zhang X, Iskander SM and Smith AL (2021) Solid waste: an overlooked source of microplastics to the environment. *Science of the Total Environment* **769**: article 144581, <https://doi.org/10.1016/j.scitotenv.2020.144581>.
- Griffin DW (2007) Atmospheric movement of microorganisms in clouds of desert dust and implications for human health. *Clinical Microbiology Reviews* **20**(3): 459–477, <https://doi.org/10.1128/CMR.00039-06>.
- Grigoratos T and Martini G (2014) *Non-exhaust Traffic Related Emissions – Brake and Tyre Wear PM*. Publications Office of the European Union, Luxembourg, Luxembourg.
- Guerranti C, Martellini T, Perra G, Scopetani C and Cincinelli A (2019) Microplastics in cosmetics: environmental issues and needs for global bans. *Environmental Toxicology and Pharmacology* **68**: 75–79, <https://doi.org/10.1016/j.etap.2019.03.007>.
- Guo JJ, Huang XP, Xiang L *et al.* (2020) Source, migration and toxicology of microplastics in soil. *Environment International* **137**: article 105263, <https://doi.org/10.1016/j.envint.2019.105263>.
- Hansen AL, Donnelly C, Refsgaard JC and Karlsson IB (2018) Simulation of nitrate reduction in groundwater – an upscaling approach from small catchments to the Baltic Sea basin. *Advances in Water Resources* **111**: 58–69, <https://doi.org/10.1016/j.advwatres.2017.10.024>.
- Harrison RM, Jones AM, Gietl J, Yin J and Green DC (2012) Estimation of the contributions of brake dust, tire wear, and resuspension to nonexhaust traffic particles derived from atmospheric measurements.

Offprint provided courtesy of www.icevirtuallibrary.com
 Author copy for personal use, not for distribution

- Environmental Science and Technology* **46**(12): 6523–6529, <https://doi.org/10.1021/es300894r>.
- Hartmann NB, Hüffer T, Thompson RC *et al.* (2019) Are we speaking the same language? Recommendations for a definition and categorization framework for plastic debris. *Environmental Science and Technology* **53**(3): 1039–1047, <https://doi.org/10.1021/acs.est.8b05297>.
- He P, Chen L, Shao L, Zhang H and Lü F (2019) Municipal solid waste (MSW) landfill: a source of microplastics? – Evidence of microplastics in landfill leachate. *Water Research* **159**: 38–45, <https://doi.org/10.1016/j.watres.2019.04.060>.
- Helmlinger MS, Tiemann LK and Grieshop MJ (2020) Towards an ecology of soil microplastics. *Functional Ecology* **34**(3): 550–560, <https://doi.org/10.1111/1365-2435.13495>.
- Horton AA, Walton A, Spurgeon DJ, Lahive E and Svendsen C (2017) Microplastics in freshwater and terrestrial environments: evaluating the current understanding to identify the knowledge gaps and future research priorities. *Science of the Total Environment* **586**: 127–141, <https://doi.org/10.1016/j.scitotenv.2017.01.190>.
- Huang Y, Liu Q, Jia W, Yan C and Wang J (2020) Agricultural plastic mulching as a source of microplastics in the terrestrial environment. *Environmental Pollution* **260**: article 114096, <https://doi.org/10.1016/j.envpol.2020.114096>.
- Huerta Lwanga E, Gertsens H, Gooren H *et al.* (2017) Incorporation of microplastics from litter into burrows of *Lumbricus terrestris*. *Environmental Pollution* **220**(Part A): 523–531, <https://doi.org/10.1016/j.envpol.2016.09.096>.
- Jahnke A, Arp HPH, Escher BI *et al.* (2017) Reducing uncertainty and confronting ignorance about the possible impacts of weathering plastic in the marine environment. *Environmental Science and Technology Letters* **4**(3): 85–90, <https://doi.org/10.1021/acs.estlett.7b00008>.
- Ji X, Ma Y, Zeng G *et al.* (2021) Transport and fate of microplastics from riverine sediment dredge piles: implications for disposal. *Journal of Hazardous Materials* **404**(Part A): article 124132, <https://doi.org/10.1016/j.jhazmat.2020.124132>.
- Jiang X, Chang Y, Zhang T *et al.* (2020) Toxicological effects of polystyrene microplastics on earthworm (*Eisenia fetida*). *Environmental Pollution* **259**: article 113896, <https://doi.org/10.1016/j.envpol.2019.113896>.
- Johnson AC, Ball H, Cross R *et al.* (2020) Identification and quantification of microplastics in potable water and their sources within water treatment works in England and Wales. *Environmental Science and Technology* **54**(19): 12326–12334, <https://doi.org/10.1021/acs.est.0c03211>.
- Kasirajan S and Ngouajio M (2012) Polyethylene and biodegradable mulches for agricultural applications: a review. *Agronomy for Sustainable Development* **32**: 501–529, <https://doi.org/10.1007/s13593-011-0068-3>.
- Katsumi N, Kusube T, Nagao S and Okochi H (2021) Accumulation of microcapsules derived from coated fertilizer in paddy fields. *Chemosphere* **267**: article 129185, <https://doi.org/10.1016/j.chemosphere.2020.129185>.
- Kedzierski M, Lechat B, Sire O *et al.* (2020) Microplastic contamination of packaged meat: occurrence and associated risks. *Food Packaging and Shelf Life* **24**: article 100489, <https://doi.org/10.1016/j.fpsl.2020.100489>.
- Kim SW and An YJ (2019) Soil microplastics inhibit the movement of springtail species. *Environment International* **126**: 699–706, <https://doi.org/10.1016/j.envint.2019.02.067>.
- Knobloch E, Ruffell H, Aves A *et al.* (2021) Comparison of deposition sampling methods to collect airborne microplastics in Christchurch, New Zealand. *Water, Air, and Soil Pollution* **232**(4): article 133, <https://doi.org/10.1007/s11270-021-05080-9>.
- Kole PJ, Löhr AJ, Van Belleghem FGAJ and Ragas AMJ (2017) Wear and tear of tyres: a stealthy source of microplastics in the environment. *International Journal of Environmental Research and Public Health* **14**(10): article 1265, <https://doi.org/10.3390/ijerph14101265>.
- Koskei K, Munyasya AN, Wang YB *et al.* (2021) Effects of increased plastic film residues on soil properties and crop productivity in agro-ecosystem. *Journal of Hazardous Materials* **414**: article 125521, <https://doi.org/10.1016/j.jhazmat.2021.125521>.
- Koutnik VS, Alkidim S, Leonard J *et al.* (2021) Unaccounted microplastics in wastewater sludge: where do they go? *ACS ES&T Water* **1**(5): 1086–1097, <https://doi.org/10.1021/acsestwater.0c00267>.
- Kwak JI and An YJ (2021) Microplastic digestion generates fragmented nanoplastics in soils and damages earthworm spermatogenesis and coelomocyte viability. *Journal of Hazardous Materials* **402**: article 124034, <https://doi.org/10.1016/j.jhazmat.2020.124034>.
- Lackmann C, Velki M, Šimić A *et al.* (2022) Two types of microplastics (polystyrene-HBCD and car tire abrasion) affect oxidative stress-related biomarkers in earthworm *Eisenia andrei* in a time-dependent manner. *Environment International* **163**: article 107190, <https://doi.org/10.1016/j.envint.2022.107190>.
- Lear G, Kingsbury JM, Franchini S *et al.* (2021) Plastics and the microbiome: impacts and solutions. *Environmental Microbiome* **16**: article 2, <https://doi.org/10.1186/s40793-020-00371-w>.
- Leslie HA, van Velzen MJM, Brandsma SH *et al.* (2022) Discovery and quantification of plastic particle pollution in human blood. *Environmental International* **163**: article 107199, <https://doi.org/10.1016/j.envint.2022.107199>.
- Li C, Moore-Kucera J, Lee J *et al.* (2014) Effects of biodegradable mulch on soil quality. *Applied Soil Ecology* **79**: 59–69, <https://doi.org/10.1016/j.apsoil.2014.02.012>.
- Li Z, Li Q, Li R, Zhou J and Wang G (2021) The distribution and impact of polystyrene nanoplastics on cucumber plants. *Environmental Science and Pollution Research* **28**(13): 16042–16053, <https://doi.org/10.1007/s11356-020-11702-2>.
- Liu K, Wang X, Song Z, Wei N and Li D (2020) Terrestrial plants as a potential temporary sink of atmospheric microplastics during transport. *Science of the Total Environment* **742**: article 140523, <https://doi.org/10.1016/j.scitotenv.2020.140523>.
- Loppi S, Roblin B, Paoli L and Aherne J (2021) Accumulation of airborne microplastics in lichens from a landfill dumping site (Italy). *Scientific Reports* **11**: article 4564, <https://doi.org/10.1038/s41598-021-84251-4>.
- Luo Y, Zhang Y, Xu Y, Guo X and Zhu L (2020) Distribution characteristics and mechanism of microplastics mediated by soil physicochemical properties. *Science of the Total Environment* **726**: article 138389, <https://doi.org/10.1016/j.scitotenv.2020.138389>.
- Luo Y, Li L, Feng Y *et al.* (2022) Quantitative tracing of uptake and transport of submicrometre plastics in crop plants using lanthanide chelates as a dual-functional tracer. *Nature Nanotechnology* **17**: 424–431, <https://doi.org/10.1038/s41565-021-01063-3>.
- Lwanga EH, Gertsens H, Gooren H *et al.* (2016) Microplastics in the terrestrial ecosystem: implications for *Lumbricus terrestris* (Oligochaeta, Lumbricidae). *Environmental Science and Technology* **50**(5): 2685–2691, <https://doi.org/10.1021/acs.est.5b05478>.
- Lwanga EH, Beriot N, Corradini F *et al.* (2022) Review of microplastic sources, transport pathways and correlations with other soil stressors: a journey from agricultural sites into the environment. *Chemical and Biological Technologies in Agriculture* **9**: article 20, <https://doi.org/10.1186/s40538-021-00278-9>.
- Mahon AM, O'Connell B, Healy MG *et al.* (2017) Microplastics in sewage sludge: effects of treatment. *Environmental Science and Technology* **51**(2): 810–818, <https://doi.org/10.1021/acs.est.6b04048>.
- Manikanda Bharath K, Usha N, Vaikunth R *et al.* (2021) Spatial distribution of microplastic concentration around landfill sites and its potential risk on groundwater. *Chemosphere* **277**: article 130263, <https://doi.org/10.1016/j.chemosphere.2021.130263>.
- Mateos-Cárdenas A, Scott DT, Seitmaganbetova G *et al.* (2019) Polyethylene microplastics adhere to *Lemna minor* (L.), yet have no

Offprint provided courtesy of www.icevirtuallibrary.com
 Author copy for personal use, not for distribution

- effects on plant growth or feeding by *Gammarus duebeni* (Lillj.). *Science of the Total Environment* **689**: 413–421, <https://doi.org/10.1016/j.scitotenv.2019.06.359>.
- Mateos-Cárdenas A, van Pelt FNAM, O'Halloran J and Jansen MAK (2021) Adsorption, uptake and toxicity of micro- and nanoplastics: effects on terrestrial plants and aquatic macrophytes. *Environmental Pollution* **284**: article 117183, <https://doi.org/10.1016/j.envpol.2021.117183>.
- Mattsson K, Johnson EV, Malmendal A *et al.* (2017) Brain damage and behavioural disorders in fish induced by plastic nanoparticles delivered through the food chain. *Scientific Reports* **7**(1): article 11452, <https://doi.org/10.1038/s41598-017-10813-0>.
- Mishra S, Rath CC and Das AP (2019) Marine microfiber pollution: a review on present status and future challenges. *Marine Pollution Bulletin* **140**: 188–197, <https://doi.org/10.1016/j.marpolbul.2019.01.039>.
- Mohamed Nor NH, Kooi M, Diepens NJ and Koelmans AA (2021) Lifetime accumulation of microplastic in children and adults. *Environmental Science and Technology* **55**(8): 5084–5096, <https://doi.org/10.1021/acs.est.0c07384>.
- Monkul MM and Özhan HO (2021) Microplastic contamination in soils: a review from geotechnical engineering view. *Polymers* **13**: article 4129, <https://doi.org/10.3390/polym13234129>.
- Muroi F, Tachibana Y, Kobayashi Y, Sakurai T and Kasuya KI (2016) Influences of poly(butylene adipate-co-terephthalate) on soil microbiota and plant growth. *Polymer Degradation and Stability* **129**: 338–346, <https://doi.org/10.1016/j.polymdegradstab.2016.05.018>.
- Ng EL, Lwanga EH, Eldridge SM *et al.* (2018) An overview of microplastic and nanoplastic pollution in agroecosystems. *Science of the Total Environment* **627**: 1377–1388, <https://doi.org/10.1016/j.scitotenv.2018.01.341>.
- Ng EL, Lin SY, Dungan AM *et al.* (2021) Microplastic pollution alters forest soil microbiome. *Journal of Hazardous Materials* **409**: article 124606, <https://doi.org/10.1016/j.jhazmat.2020.124606>.
- Nizzetto L, Futter M and Langaas S (2016) Are agricultural soils dumps for microplastics of urban origin? *Environmental Science and Technology* **50**(20): 10777–10779, <https://doi.org/10.1021/acs.est.6b04140>.
- O'Connor D, Pan S, Shen Z *et al.* (2019) Microplastics undergo accelerated vertical migration in sand soil due to small size and wet–dry cycles. *Environmental Pollution* **249**: 527–534, <https://doi.org/10.1016/j.envpol.2019.03.092>.
- O'Kelly BC and Soltani A (2022a) Discussion of “Behaviour of a foam mixture as a lightweight construction material” [Int J of Geosynth and Ground Eng (2021) 7(3), 51]. *International Journal of Geosynthetics and Ground Engineering* **8**(2): article 30, <https://doi.org/10.1007/s40891-022-00369-z>.
- O'Kelly BC and Soltani A (2022b) Discussion: Effects of plastic waste materials on geotechnical properties of clayey soil [DOI: 10.1007/s40515-020-00145-4]. *Transportation Infrastructure Geotechnology*, <https://doi.org/10.1007/s40515-022-00224-8>.
- O'Kelly BC, El-Zein A, Liu X *et al.* (2021) Microplastics in soils: an environmental geotechnics perspective. *Environmental Geotechnics* **8**(8): 586–618, <https://doi.org/10.1680/jenge.20.00179>.
- Oliveri Conti G, Ferrante M, Banni M *et al.* (2020) Micro- and nanoplastics in edible fruit and vegetables. The first diet risks assessment for the general population. *Environmental Research* **187**: article 109677, <https://doi.org/10.1016/j.envres.2020.109677>.
- Otake Y, Kobayashi T, Asabe H, Murakami N and Ono K (1995) Biodegradation of low-density polyethylene, polystyrene, polyvinyl chloride, and urea formaldehyde resin buried under soil for over 32 years. *Applied Polymer Science* **56**(13): 1789–1796, <https://doi.org/10.1002/app.1995.070561309>.
- Panno SV, Kelly WR, Scott J *et al.* (2019) Microplastic contamination in karst groundwater systems. *Groundwater* **57**(2): 189–196, <https://doi.org/10.1111/gwat.12862>.
- Pauly JL, Stegmeier SJ, Allaart HA *et al.* (1998) Inhaled cellulosic and plastic fibers found in human lung tissue. *Cancer Epidemiology, Biomarkers & Prevention* **7**(5): 419–428.
- Powell JJ, Faria N, Thomas-McKay E and Pele LC (2010) Origin and fate of dietary nanoparticles and microparticles in the gastrointestinal tract. *Journal of Autoimmunity* **34**(3): 226–233, <https://doi.org/10.1016/j.jaut.2009.11.006>.
- Prata JC (2018) Airborne microplastics: consequences to human health? *Environmental Pollution* **234**: 115–126, <https://doi.org/10.1016/j.envpol.2017.11.043>.
- Qi Y, Yang X, Pelaez AM *et al.* (2018) Macro- and micro- plastics in soil–plant system: effects of plastic mulch film residues on wheat (*Triticum aestivum*) growth. *Science of the Total Environment* **645**: 1048–1056, <https://doi.org/10.1016/j.scitotenv.2018.07.229>.
- Qi R, Jones DL, Li Z, Liu Q and Yan C (2020a) Behavior of microplastics and plastic film residues in the soil environment: a critical review. *Science of the Total Environment* **703**: article 134722, <https://doi.org/10.1016/j.scitotenv.2019.134722>.
- Qi Y, Ossowicki A, Yang X *et al.* (2020b) Effects of plastic mulch film residues on wheat rhizosphere and soil properties. *Journal of Hazardous Materials* **387**: article 121711, <https://doi.org/10.1016/j.jhazmat.2019.121711>.
- Ragusa A, Svelato A, Santacroce C *et al.* (2021) Plasticenta: first evidence of microplastics in human placenta. *Environment International* **146**: article 106274, <https://doi.org/10.1016/j.envint.2020.106274>.
- Rasmussen LA, Iordachescu L, Tumlin S and Vollertsen J (2021) A complete mass balance for plastics in a wastewater treatment plant – microplastics contributes more than microplastics. **201**: article 117307, <https://doi.org/10.1016/j.watres.2021.117307>.
- Ren Z, Gui X, Xu X *et al.* (2021) Microplastics in the soil–groundwater environment: aging, migration, and co-transport of contaminants – a critical review. *Journal of Hazardous Materials* **419**: article 126455, <https://doi.org/10.1016/j.jhazmat.2021.126455>.
- Ribeiro F, O'Brien JW, Galloway T and Thomas KV (2019) Accumulation and fate of nano- and micro-plastics and associated contaminants in organisms. *TrAC Trends in Analytical Chemistry* **111**: 139–147, <https://doi.org/10.1016/j.trac.2018.12.010>.
- Rillig MC, Ingrassia R and de Souza Machado AA (2017a) Microplastic incorporation into soil in agroecosystems. *Frontiers in Plant Science* **8**: article 1805, <https://doi.org/10.3389/fpls.2017.01805>.
- Rillig MC, Ziersch L and Hempel S (2017b) Microplastic transport in soil by earthworms. *Scientific Reports* **7**: article 1362, <https://doi.org/10.1038/s41598-017-01594-7>.
- Rodland ES, Samanipour S, Rauert C *et al.* (2022) A novel method for the quantification of tire and polymer-modified bitumen particles in environmental samples by pyrolysis gas chromatography mass spectroscopy. *Journal of Hazardous Materials* **423**: article 127092, <https://doi.org/10.1016/j.jhazmat.2021.127092>.
- Ruffell H, Pantos O, Northcott G and Gaw S (2021) Wastewater treatment plant effluents in New Zealand are a significant source of microplastics to the environment. *New Zealand Journal of Marine and Freshwater Research*, <https://doi.org/10.1080/00288330.2021.1988647>.
- Samandra S, Johnston JM, Jaeger JE *et al.* (2022) Microplastic contamination of an unconfined groundwater aquifer in Victoria, Australia. *Science of the Total Environment* **802**: article 149727, <https://doi.org/10.1016/j.scitotenv.2021.149727>.
- Schneider M, Stracke F, Hansen S and Schaefer UF (2009) Nanoparticles and their interactions with the dermal barrier. *Dermato-Endocrinology* **1**(4): 197–206, <https://doi.org/10.4161/derm.1.4.9501>.
- Schwabl P, Köppel S, Königshofer P *et al.* (2019) Detection of various microplastics in human stool – a prospective case series. *Annals of Internal Medicine* **171**(7): 453–457, <https://doi.org/10.7326/M19-0618>.
- Schwab F, Rothen-Rutishauser B and Petri-Fink A (2020) When plants and plastic interact. *Nature Nanotechnology* **15**: 729–730, <https://doi.org/10.1038/s41565-020-0762-x>.

Offprint provided courtesy of www.icevirtuallibrary.com
 Author copy for personal use, not for distribution

- Schwarz AE, Ligthart TN, Boukris E and van Harmelen T (2019) Sources, transport, and accumulation of different types of plastic litter in aquatic environments: a review study. *Marine Pollution Bulletin* **143**: 92–100, <https://doi.org/10.1016/j.marpolbul.2019.04.029>.
- Selvam S, Jesuraja K, Venkatramanan S, Roy PD and Kumari VJ (2021) Hazardous microplastic characteristics and its role as a vector of heavy metal in groundwater and surface water of coastal South India. *Journal of Hazardous Materials* **402**: article 123786, <https://doi.org/10.1016/j.jhazmat.2020.123786>.
- Senathirajah K, Attwood S, Bhagwat G *et al.* (2021) Estimation of the mass of microplastics ingested – a pivotal first step towards human health risk assessment. *Journal of Hazardous Materials* **404(Part B)**: article 124004, <https://doi.org/10.1016/j.jhazmat.2020.124004>.
- Shamskhany A, Li Z, Patel P and Karimpour S (2021) Evidence of microplastic size impact on mobility and transport in the marine environment: a review and synthesis of recent research. *Frontiers in Marine Science* **8**: article 760649, <https://doi.org/10.3389/fmars.2021.760649>.
- Shi J, Wu D, Su Y and Xie B (2021) Selective enrichment of antibiotic resistance genes and pathogens on polystyrene microplastics in landfill leachate. *Science of the Total Environment* **765**: article 142775, <https://doi.org/10.1016/j.scitotenv.2020.142775>.
- Sobhani Z, Lei Y, Tang Y *et al.* (2020) Microplastics generated when opening plastic packaging. *Scientific Reports* **10**: article 4841, <https://doi.org/10.1038/s41598-020-61146-4>.
- Šourková M, Adamcová D, Winkler J and Vavřková MD (2021a) Phytotoxicity of tires evaluated in simulated conditions. *Environments* **8(6)**: article 49, <https://doi.org/10.3390/environments8060049>.
- Šourková M, Adamcová D and Vavřková MD (2021b) The influence of microplastics from ground tyres on the acute, subchronical toxicity and microbial respiration of soil. *Environments* **8(11)**: article 128, <https://doi.org/10.3390/environments8110128>.
- Steinmetz Z, Wollmann C, Schaefer M *et al.* (2016) Plastic mulching in agriculture. Trading short-term agronomic benefits for long-term soil degradation? *Science of the Total Environment* **550**: 690–705, <https://doi.org/10.1016/j.scitotenv.2016.01.153>.
- Su Y, Zhang Z, Wu D *et al.* (2019) Occurrence of microplastics in landfill systems and their fate with landfill age. *Water Research* **164**: article 114968, <https://doi.org/10.1016/j.watres.2019.114968>.
- Su Y, Zhang Z, Zhu J *et al.* (2021) Microplastics act as vectors for antibiotic resistance genes in landfill leachate: the enhanced roles of the long-term aging process. *Environmental Pollution* **270**: article 116278, <https://doi.org/10.1016/j.envpol.2020.116278>.
- Taltec K, Huvet A, Yeuc'h V, Le Goïc N and Paul-Pont I (2022) Chemical effects of different types of rubber-based products on early life stages of Pacific oyster, *Crassostrea gigas*. *Journal of Hazardous Materials* **427**: article 127883, <https://doi.org/10.1016/j.jhazmat.2021.127883>.
- Thompson RC, Olsen Y, Mitchell RP *et al.* (2004) Lost at sea: where is all the plastic? *Science* **304(5672)**: 838, <https://doi.org/10.1126/science.1094559>.
- Turcotte SE, Chee A, Walsh R *et al.* (2013) Flock worker's lung disease: natural history of cases and exposed workers in Kingston, Ontario. *Chest* **143(6)**: 1642–1648, <https://doi.org/10.1378/chest.12-0920>.
- Vavřková MD and O'Kelly BC (2022) Comment on “A new approach to stabilization of calcareous dune sand” [Int. J. Environ. Sci. Technol. **19**, 3581–3592 (2022)]. *International Journal of Environmental Science and Technology*, <https://doi.org/10.1007/s13762-022-04291-9>.
- Vavřková MD, Paleologos EK, Dominijanni A *et al.* (2021) Municipal solid waste management under COVID-19: challenges and recommendations. *Environmental Geotechnics* **8(3)**: 217–232, <https://doi.org/10.1680/jenge.20.00082>.
- Viaroli S, Lancia M and Re V (2022) Microplastics contamination of groundwater: current evidence and future perspectives. A review. *Science of the Total Environment* **824**: article 153851, <https://doi.org/10.1016/j.scitotenv.2022.153851>.
- Wang YP, Li XG, Fu T *et al.* (2016) Multi-site assessment of the effects of plastic-film mulch on the soil organic carbon balance in semiarid areas of China. *Agricultural and Forest Meteorology* **228–229**: 42–51, <https://doi.org/10.1016/j.agrformet.2016.06.016>.
- Wang W, Ge J, Yu X and Li H (2020) Environmental fate and impacts of microplastics in soil ecosystems: progress and perspective. *Science of the Total Environment* **708**: article 134841, <https://doi.org/10.1016/j.scitotenv.2019.134841>.
- Wang F, Wang X and Song N (2021) Polyethylene microplastics increase cadmium uptake in lettuce (*Lactuca sativa* L.) by altering the soil microenvironment. *Science of the Total Environment* **784**: article 147133, <https://doi.org/10.1016/j.scitotenv.2021.147133>.
- Wang Y, Wang F, Xiang L *et al.* (2022) Attachment of positively and negatively charged submicron polystyrene plastics on nine typical soils. *Journal of Hazardous Materials* **431**: article 128566, <https://doi.org/10.1016/j.jhazmat.2022.128566>.
- Watteau F, Dignac MF, Bouchard A, Revallier A and Houot S (2018) Microplastics detection in soil amended with municipal solid waste composts as revealed by transmission electron microscopy and pyrolysis/GC/MS. *Frontiers in Sustainable Food Systems* **2**: article 81, <https://doi.org/10.3389/fsufs.2018.00081>.
- Wei R and Zimmermann W (2017) Microbial enzymes for the recycling of recalcitrant petroleum-based plastics: how far are we? *Microbial Biotechnology* **10(6)**: 1308–1322, <https://doi.org/10.1111/1751-7915.12710>.
- Weithmann N, Möller JN, Löder MGJ *et al.* (2018) Organic fertilizer as a vehicle for the entry of microplastic into the environment. *Science Advances* **4(4)**: article eaap8060, <https://doi.org/10.1126/sciadv.aap8060>.
- Yang Y, Vance M, Tou F *et al.* (2016) Nanoparticles in road dust from impervious urban surfaces: distribution, identification, and environmental implications. *Environmental Science: Nano* **3(3)**: 534–544, <https://doi.org/10.1039/c6en00056h>.
- Zhang M, Zhao Y, Qin X *et al.* (2019a) Microplastics from mulching film is a distinct habitat for bacteria in farmland soil. *Science of the Total Environment* **688**: 470–478, <https://doi.org/10.1016/j.scitotenv.2019.06.108>.
- Zhang Y, Gao T, Kang S and Sillanpää M (2019b) Importance of atmospheric transport for microplastics deposited in remote areas. *Environmental Pollution* **254(Part A)**: article 112953, <https://doi.org/10.1016/j.envpol.2019.07.121>.
- Zhang J, Wang L and Kannan K (2020) Microplastics in house dust from 12 countries and associated human exposure. *Environment International* **134**: article 105314, <https://doi.org/10.1016/j.envint.2019.105314>.
- Zhang X, Chen Y, Li X *et al.* (2022) Size/shape-dependent migration of microplastics in agricultural soil under simulative and natural rainfall. *Science of the Total Environment* **815**: article 152507, <https://doi.org/10.1016/j.scitotenv.2021.152507>.
- Zhu D, Bi QF, Xiang Q *et al.* (2018) Trophic predator–prey relationships promote transport of microplastics compared with the single *Hypoaspis aculeifer* and *Folsomia candida*. *Environmental Pollution* **235**: 150–154, <https://doi.org/10.1016/j.envpol.2017.12.058>.

How can you contribute?

To discuss this paper, please submit up to 500 words to the editor at journals@ice.org.uk. Your contribution will be forwarded to the author(s) for a reply and, if considered appropriate by the editorial board, it will be published as a discussion in a future issue of the journal.